

Towards a tomographic picture of a Bose-Einstein condensate

Stefano Mancini

Istituto Nazionale per la Fisica della Materia,

Dipartimento di Fisica, Università di Milano, Via Celoria 16, I-20133 Milano, Italy

Mauro Fortunato

Istituto Nazionale per la Fisica della Materia,

Dipartimento di Matematica e Fisica, Università di Camerino, I-62032 Camerino, Italy

Paolo Tombesi

Istituto Nazionale per la Fisica della Materia,

Dipartimento di Matematica e Fisica, Università di Camerino, I-62032 Camerino, Italy

Giacomo Mauro D'Ariano

Istituto Nazionale per la Fisica della Materia,

Dipartimento di Fisica "A. Volta", Università di Pavia, I-27100 Pavia, Italy

(Received: February 1, 2008)

Abstract

We investigate by means of numerical simulations the possibilities of tomographic techniques applied to a Bose-Einstein condensate in order to reconstruct its ground state. Essentially, two scenarios are considered for which the density matrix elements can be retrieved from atom counting probabilities. The methods presented here allow to distinguish among various possible quantum states.

OCIS key terms: Quantum Optics 270.0270, Atomic and molecular physics

020.0020

PACS number(s): 03.65.Bz, 03.75.Fi, 32.80.-t

Typeset using REVTeX

I. INTRODUCTION

Before the birth of quantum mechanics, optics and mechanics have long developed on parallel tracks, as light and massive entities were considered as waves and particles, respectively. At the beginning of the 20th century, with the introduction of quantum mechanics, waves and particles started to play an interchangeable role, with the concepts of photons and of De Broglie wavelength. This gave rise to the birth of quantum optics and atom optics [1]. However, while an optical single-mode system has already been available since long ago, the same cannot be said for matter waves. In fact, in the field of atom optics, only recently breakthroughs in the evaporative cooling of dilute alkali gases have allowed the generation of Bose-Einstein condensates (BEC) [2]. The BEC is a macroscopic occupation of the ground state of the gas and is one important paradigm of quantum statistical mechanics.

In recent theoretical and experimental investigations of BEC, one of the most important and urgent issues is the determination of the actual quantum state of the condensate. In fact, as in optics, the presence of many particles in a single mode makes it possible to inquire about the multiparticle quantum state of the mode. At first thought, a number state might seem a natural description of a condensate mode, but the actual state may well depend on the details of preparation. For example the demonstration of first-order interference [3] and observations of normalized, spatial correlation functions [4] near unity suggest coherent states, while the presence of collisions between atoms may lead to the formation of squeezed states [5]. Also the internal states of the condensate atoms allow precise manipulation of the BEC state by interaction with light [6].

Few experimental methods of obtaining partial information about the state have been suggested [7]. On the other hand, motivated by the success of quantum tomographic techniques in optics [8], more direct methods for measuring the quantum state of a BEC were recently proposed [9–11].

In optical tomography, the key point is the use of a reference field, namely the local oscillator [8]. The latter, prepared in a coherent state, allows one to probe the desired state

through the measurement of a set of probabilities [12]. However, a difficulty arises if we try to adapt the same technique to a BEC. In fact, while in optics it is easy to obtain a coherent reference field (e.g. from a laser), the same is not actually available for atoms. Nevertheless, recent progresses in this direction seem promising [13,14]. Hence, in the present paper, we provide a detailed study of the possibility to reconstruct the quantum state of a BEC which includes both the scenarios: when a reference field is available and when it is not.

The paper is organized as follows: in Section II we review the tomographic principle and consider a suitable operator transform on the atomic system. In Section III, we consider the case of state reconstruction in the absence of a reference field, while in Section IV the opposite case is analyzed. Finally, in Section V we comment the numerical results, and in Section VI we conclude with a brief discussion.

II. THE BASIC MODEL

Recently, we established [15] a quite general principle of constructing measurable probabilities, which determine completely the quantum state in the tomographic approach. For a more refined treatment see Ref. [16].

Let us consider a quantum state described by the density operator $\hat{\rho}$, which is a nonnegative Hermitian operator, i.e.,

$$\hat{\rho}^\dagger = \hat{\rho}, \quad \text{Tr } \hat{\rho} = 1, \quad (1)$$

and

$$\langle v | \hat{\rho} | v \rangle = \rho_{v,v} \geq 0. \quad (2)$$

We label the vector basis $|v\rangle$ in the space of pure quantum states by the index v which may represent any degrees of freedom of the system under consideration. Formula (2) can be rewritten by using the Hermitian projection operator

$$\hat{\Pi}_v = |v\rangle\langle v|, \quad (3)$$

in the following form

$$\rho_{v,v} = \text{Tr} \left(\hat{\Pi}_v \hat{\rho} \right) . \quad (4)$$

On the other hand, in the space of states, there will be a family of unitary transformation operators $\hat{U}(\sigma)$ depending on some parameters $\sigma = (\sigma_1, \dots, \sigma_k \dots)$, that can be sometimes identified with a group-representation operators. It was shown [15] that known tomography schemes can be considered from the viewpoint of group theory by using appropriate groups. More recently this concept has been developed obtaining an elegant group theoretical approach to quantum state measurement [16]. Here, we formulate the tomographic approach in the following way. Let us introduce a “transformed density operator”

$$\hat{\rho}_\sigma = \hat{U}^{-1}(\sigma) \hat{\rho} \hat{U}(\sigma) . \quad (5)$$

Its diagonal elements are still nonnegative probabilities

$$\langle v | \hat{\rho}_\sigma | v \rangle \equiv w(v, \sigma) \geq 0 . \quad (6)$$

These probabilities are functions of stochastic variable(s) v and parameter(s) σ . As a consequence of the unit trace of the density operator, they also fulfill the normalization condition

$$\int dv w(v, \sigma) = 1 . \quad (7)$$

Of course, in the case of discrete indices, the integral in Eq. (7) should be replaced by a sum over discrete variables.

The l.h.s. of Eq. (6) can be interpreted as the probability density for the measurement of the observable $\hat{V}$ (the operator whose eigenstates are given by $|v\rangle$) in an ensemble of transformed reference frames labeled by the index σ , if the state $\hat{\rho}$ is given. Along with this interpretation, one can also consider the transformed projector

$$\hat{\Pi}_v(\sigma) = \hat{U}(\sigma) \hat{\Pi}_v \hat{U}^{-1}(\sigma) , \quad (8)$$

in terms of which the expression (6) for the probability $w(v, \sigma)$ takes the form

$$w(v, \sigma) = \text{Tr} [\hat{\rho} \hat{\Pi}_v(\sigma)] . \quad (9)$$

These probability densities are also called “marginal” distributions as a generalization of the concept introduced by Wigner [17]. The tomography schemes are based on the possibility to find the inverse of Eq. (9). If it is possible to solve Eq. (9), considering the probability $w(v, \sigma)$ as a known function and the density matrix as an unknown operator, the quantum state can be reconstructed in terms of measurable positive definite probability distributions. This is the essence of state reconstruction techniques.

Specifically, we consider two atomic sources whose atoms (described by two bosonic modes $\hat{b}_1$ and $\hat{b}_2$) can be mixed through an atomic beam splitter [18], and assume that successively a phase shift ϕ can also be introduced between them. We shall specify these modes later. At the output, a detection of the number of atoms in both modes can be performed. This amounts to the possibility of measuring the probability distributions related to the transformed state

$$\hat{\rho} \rightarrow \hat{U}(\theta, \phi) \hat{\rho} \hat{U}^\dagger(\theta, \phi) , \quad (10)$$

where the transformation operator is given by

$$\hat{U}(\theta, \phi) = \exp \left\{ -i \frac{\theta}{2} \left[\hat{b}_1^\dagger \hat{b}_2 \exp(-i\phi) + \hat{b}_1 \hat{b}_2^\dagger \exp(i\phi) \right] \right\} . \quad (11)$$

Here, $\cos^2(\theta/2)$ represents the transmission coefficient at the beam splitter. Eq. (10) plays the same role of Eq. (5), and, in the spirit of the tomographic principle, the set of “marginals” associated to the transformed state will allow us to recover the original state. In the next two Sections, as anticipated in the Introduction, we shall distinguish two situations.

III. CASE I

We first treat the case where a reference field is not available. All we can do in this case is to consider two condensates belonging to the two modes $\hat{b}_1$, $\hat{b}_2$, and put the constraint of

total particle number conservation, i.e. $[\hat{\rho}, \hat{N}] = 0$, in order to infer their (joint) state. The latter is assumed to be a generic two-mode state of the type

$$|\Psi\rangle = \sum_{n=0}^N c_n |N-n\rangle_1 |n\rangle_2. \quad (12)$$

At this stage we use the formal equivalence between the algebra for two harmonic oscillators and that for angular momentum [19]. We write the state $|n\rangle_1 |N-m\rangle_2 = |j+m\rangle_1 |j-m\rangle_2$ as a spin state $|m\rangle$, where $j = N/2$ and $m = n - j$ ($m = -j, -j+1, \dots, j-1, j$). The $j+1$ states $|m\rangle$ have all the properties of the eigenstates of $\hat{J}^2$ and $\hat{J}_z$ where

$$\hat{J}_+ = \hat{J}_-^\dagger = \hat{b}_1^\dagger \hat{b}_2, \quad \hat{J}_z = \frac{1}{2} (\hat{b}_1^\dagger \hat{b}_1 - \hat{b}_2^\dagger \hat{b}_2), \quad \hat{J}^2 = \hat{J}_z^2 + \frac{1}{2} (\hat{J}_+ \hat{J}_- + \hat{J}_- \hat{J}_+). \quad (13)$$

The effect of the beam splitter, including the phase shift, is a rotation by an angle θ about an axis $\mathbf{u}_\phi = \mathbf{u}_x \cos \phi - \mathbf{u}_y \sin \phi$ of the angular momenta $\hat{\mathbf{J}}$, i.e.

$$\hat{U}(\theta, \phi) = \exp(-i\theta \hat{\mathbf{J}} \cdot \mathbf{u}_\phi). \quad (14)$$

On the other hand, the rotation (14) can be specified by means of the Wigner-D function [19]

$$\langle m' | \hat{U}(\theta, \phi) | m \rangle \equiv \mathcal{D}_{m' m}^{(j)}(\psi = 0, \theta, \phi), \quad (15)$$

where now ψ, θ, ϕ represent the Euler's angles. Then, the probability of $j+m$ counts at the first detector and $j-m$ at the second one is given by

$$w(m, \theta, \phi) = \sum_{m_1=-j}^j \sum_{m_2=-j}^j \mathcal{D}_{m m_1}^{(j)}(\psi, \theta, \phi) \rho_{m_1 m_2}^{(j)} \mathcal{D}_{m m_2}^{(j)*}(\psi, \theta, \phi). \quad (16)$$

The measurement of the atomic number in both modes guarantees a unit efficiency. In fact, data for which the sum of counts is not N can be disregarded. Moreover, in Eq. (16) we have left the argument ψ unspecified in the r.h.s. and omitted it in the l.h.s. since $\mathcal{D}_{m m'}^{(j)} \propto \exp(-im\psi)$: the marginal distribution only depends on the two angles θ and ϕ .

Following [20,21] we will derive the expression for the density matrix of a spin state in terms of measurable probability distributions. This can be done by using the known integral

product of three Wigner-D functions over the rotation group and the orthogonality of the Wigner- $3j$ symbols $\mathcal{W}_{m_1 m_2 m_3}^{j_1 j_2 j_3}$ [19]. Finally, the density matrix elements can be expressed in terms of the marginal distribution as

$$\begin{aligned} \rho_{m_1 m_2}^{(j)} &= (-1)^{m_2} \sum_{j'=0}^{2j} \sum_{m'=-j'}^{j'} (2j'+1)^2 \sum_{m=-j}^j \int (-1)^m w(m, \theta, \phi) \\ &\times \mathcal{D}_{0 m'}^{(j')}(\psi, \theta, \phi) \mathcal{W}_{m-m_2 0}^{j j'} \mathcal{W}_{m_1 -m_2 m'}^{j j'} \frac{d\Omega}{8\pi^2} \end{aligned} \quad (17)$$

where the integration is performed over the rotation parameters, i.e.

$$\int d\Omega = \int_0^{2\pi} d\psi \int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\phi. \quad (18)$$

Thus, Eq. (17) can be used to sample two-mode BEC density matrix elements starting from the measurable probabilities $w(m, \theta, \phi)$ and some known functions.

IV. CASE II

Recent progress in the generation of an atomic coherent source [13] makes us hope about the possibility to have an atomic reference field [14]. Thus, we shall consider in this section the first mode as the condensate to be investigated, and the second one coming from a coherent atomic source.

Let $\hat{\rho}$ be the state of mode 1 we want to reconstruct and $\bar{\beta}$ the coherent state characterizing mode 2. Then, the probability of counting n atoms in mode 1 for $\theta = \pi/2$, will be

$$w(n, \beta) = \text{Tr} \left[\hat{U}^{-1}(\theta = \pi/2, \phi) \hat{\rho} |\bar{\beta}\rangle_{22} \langle \bar{\beta}| \hat{U}(\theta = \pi/2, \phi) |n\rangle_{11} \langle n| \right], \quad (19)$$

where $\beta = |\bar{\beta}| \exp(i\varphi)$, $\varphi = \arg \bar{\beta} - \phi + \pi/2$. This corresponds to the probability distribution for the measurement of the displaced number operator $\hat{D}^\dagger(\beta) \hat{b}_1^\dagger \hat{b}_1 \hat{D}(\beta)$ analogously to the Photon Number Tomography [22]. In that case, however, one has to collect number distributions by spanning the whole complex plane β ; here, instead, we will simplify the procedure (see also [23]).

Of course, the number of atoms in the condensate, though not fixed, will be finite, thus, it happens that $\langle k|\rho|m\rangle = 0$ for $k, m > N_1$, with N_1 a suitable estimation of the maximum number of the condensed atoms. By virtue of this assumption we can rewrite Eq. (19) as

$$w(n, \beta) = \exp(-|\beta|^2) n! \sum_{k,m=0}^{N_1} \langle k|\rho|m\rangle \frac{1}{\sqrt{k!m!}} |\beta|^{m+k-2n} \exp[i(m-k)\varphi] L_n^{(m-n)}(|\beta|^2) \times L_n^{(k-n)}(|\beta|^2), \quad (20)$$

where $L_n^{(m)}$ are the associated Laguerre polynomials.

Let us now consider, for a given value of $|\beta|$, the function $w(n, \beta)$ as function of φ and calculate the coefficients of the Fourier expansion

$$w^{(s)}(n, |\beta|) = \frac{1}{2\pi} \int_0^{2\pi} d\varphi w(n, \beta) \exp(is\varphi), \quad (21)$$

($s = 0, 1, 2, \dots$). By combining Eqs. (20) and (21), we get

$$w^{(s)}(n, |\beta|) = \sum_{m=0}^{N_1-s} \mathcal{A}_{n,m}^{(s)}(|\beta|) \langle m+s|\rho|m\rangle, \quad (22)$$

where

$$\mathcal{A}_{n,m}^{(s)}(|\beta|) = \exp(-|\beta|^2) n! \frac{1}{\sqrt{(m+s)!m!}} |\beta|^{2(m-n)+s} L_n^{(m-n)}(|\beta|^2) L_n^{(m+s-n)}(|\beta|^2). \quad (23)$$

If the distribution $w(n, \beta)$ is measured for $n = 0, 1, \dots, N$ ($N \geq N_1$), then Eq. (22) represents for each value of s a system of $(N+1)$ linear equations between the $(N+1)$ measured quantities and the (N_1+1-s) unknown density matrix elements. Therefore, to obtain the latter we only need to invert the system [24]

$$\langle m+s|\rho|m\rangle = \sum_{n=0}^N \mathcal{M}_{m,n}^{(s)}(|\beta|) w^{(s)}(n, |\beta|), \quad (24)$$

where the matrices $\mathcal{M}$ are given by $\mathcal{M} = (\mathcal{A}^T \mathcal{A})^{-1} \mathcal{A}^T$. It is possible to see that such matrices satisfy the relation

$$\sum_{n=0}^N \mathcal{M}_{m',n}^{(s)}(|\beta|) \mathcal{A}_{n,m}^{(s)}(|\beta|) = \delta_{m,m'}, \quad (25)$$

for $m, m' = 0, 1, \dots, N_1 - s$, which means that from the exact probabilities satisfying Eq. (22) the correct density matrix is obtained. By combining Eqs. (24) and (21) we find that

$$\langle m+s|\rho|m\rangle = \frac{1}{2\pi} \sum_{n=0}^N \int d\varphi \mathcal{M}_{m,n}^{(s)}(|\beta|) \exp(is\varphi) w(n, \beta), \quad (26)$$

which may be regarded as the formula for the direct sampling of the condensate density matrix. In particular we see that the determination of the state of the condensate only requires the value of φ (i.e. the phase between reference and condensate field) to be varied. Moreover, the presented reconstruction procedure involves Laguerre polynomials in place of additional summations, which guarantee a better stability in the numerical manipulation of large set of data, with respect to analogous methods [23].

Finally, the non unit efficiency η in the detection process can be accounted for by considering a binomial convolution of the ideal probability [25]

$$w_\eta(k, \beta) = \sum_{n=k}^{\infty} \binom{n}{k} \eta^k (1-\eta)^{n-k} w(n, \beta), \quad (27)$$

and the consequent modification of the matrix $\mathcal{A}$.

V. NUMERICAL RESULTS

It is plausible, and it has been already suggested [5,11], that the state of a condensate with repulsive collisions be a squeezed state with reduced number fluctuations. Hence in the following we will consider this situation. The single 1-mode state can be written as

$$|\Psi\rangle = \sum_{n=0}^{\infty} c_n |n\rangle, \quad (28)$$

where the coefficients c_n are given by [26]

$$c_n = \left(\frac{2}{r+1}\right)^{1/2} r^{1/4} \left(\frac{r-1}{r+1}\right)^{n/2} (2^n n!)^{-1/2} H_n \left(\sqrt{\frac{2r^2}{r^2-1}} x_0 \right) \exp\left(-\frac{r}{r+1} x_0^2\right), \quad (29)$$

where r is the squeezing parameter, x_0 is the (real) displacement and H_n denotes the Hermite polynomials.

A phase space representation of this state can be given by the Q -function [27]

$$Q(\alpha) = \langle \alpha | \hat{\rho} | \alpha \rangle, \quad (30)$$

where α is the complex amplitude of a coherent state. It yields

$$Q(\alpha) = \exp(-|\alpha|^2) \left| \sum_{n=0}^{\infty} \frac{(\alpha^*)^n}{\sqrt{n!}} c_n \right|^2. \quad (31)$$

For the case discussed in Sec. III, we have to consider [5,11] a two-mode squeezed state written in the angular momentum representation, *i.e.*

$$|\Psi\rangle = \mathcal{N} \sum_{m=-j}^j c_{j+m} |m\rangle, \quad (32)$$

where $\mathcal{N}$ is a normalization factor and the coefficients c_{j+m} are given in Eq. (29). Since the quantity $x_0^2 + (r^2 - 1)/4r$ represents the mean number of atoms in mode 1, it must be smaller than the total number of atoms N . Furthermore, the atomic coherent state basis for a system of angular momentum j is defined by [28]

$$\begin{aligned} |\theta, \phi\rangle &= \sum_{m=-j}^j \mathcal{D}_{m,-j}^{(j)}(\psi=0, \theta, \phi) |m\rangle \\ &= \sum_{m=-j}^j \binom{2j}{m+j}^{1/2} \left(\sin \frac{\theta}{2} \right)^{j+m} \left(\cos \frac{\theta}{2} \right)^{j-m} \exp(-im\phi) |m\rangle, \end{aligned} \quad (33)$$

then, one can define the Q -quasiprobability distribution analogously to Eq. (30)

$$Q(\theta, \phi) = \langle \theta, \phi | \hat{\rho} | \theta, \phi \rangle. \quad (34)$$

For the state considered in Eq. (32), it becomes

$$Q(\theta, \phi) = \left| \sum_{m=-j}^j \mathcal{D}_{m,-j}^{(j)*}(\psi=0, \theta, \phi) c_{j+m} \right|^2. \quad (35)$$

This is shown in Fig. 1(a). Instead, Fig. 1(b) displays the Q -function calculated from the reconstructed density matrix elements. We may see that the method of Sec. III is quite accurate, apart from some background noise.

Analogously, in Fig. 2(a) we have plotted the ideal Wigner function [27] of Eq. (31), while Fig. 2(b) is its reconstructed version. In this case, the statistical error depends on the chosen value of $|\beta|$. For $|\beta|$ close to zero the diagonal density matrix elements can be determined very precisely, whereas the off-diagonal elements strongly fluctuate. The opposite happens

by increasing $|\beta|$. To compensate for the fluctuations, the number of measurement events must be increased. Another source of error stems from the truncation of the reconstructed density matrix at the value N_1 .

It is worth comparing the previous results with those obtained in the case of a number state. For this purpose, we show in Fig. 3(a) the ideal Wigner function for a number state. In this case, the Wigner function becomes negative, displaying the highly non classical character of a Fock state. As above, in Fig. 3(b) we show the Wigner function obtained by applying the reconstruction method of Sec. IV to such a state: the two pictures are practically indistinguishable, showing the accuracy of the present method.

Finally, as an instructive comparison we show in Fig. 4 the same Wigner function of Fig. 2(a) calculated when the state reconstruction takes place with a random phase relation between probe and condensate. As can be seen the state becomes randomized and diffused in phase, but its Wigner function remains positive. This figure should be contrasted with Fig. 3. In this case the apparent $U(1)$ symmetry is not pertaining to the state [29], but rather due to the measurement method, which implies a preparation of the (same) state at each experimental run.

VI. CONCLUSION

To conclude, we have studied, through numerical simulations, the possibilities of a tomographic approach to the quantum state of a Bose-Einstein condensate. We have considered two possible scenarios, whether an atomic reference field is available or not. The corresponding methods turn out to be accurate and robust to detection inefficiency, and allow one to distinguish among various possible quantum states of the condensate.

It is worth noting that the studied techniques allow direct sampling of the density matrix elements avoiding any ambiguities in the reconstruction procedure due to singularities [30].

The key point remains the possibility to have a reference field and/or its state preparation. Furthermore, we note the necessity to deal with a relatively small number of atoms

in order to implement efficiently the numerical algorithms. In spite of these difficulties, we retain the possibility of measuring the true density matrix of a condensate accessible and worth considering.

Finally, we would like to remark that the presented procedures could also be considered for other fields like high energy heavy ion collisions where pions can condense as well [31].

ACKNOWLEDGMENTS

The authors gratefully acknowledge discussions with R. Onofrio. This work has been partially supported by the Istituto Nazionale di Fisica della Materia under the Advanced Research Project “Cat”, the Ministero dell’Università e della Ricerca Scientifica e Tecnologica, and the European Union.

REFERENCES

- [1] D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer-Verlag, Berlin, 1994).
- [2] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, “Observation of Bose-Einstein condensation in a dilute atomic vapor”, *Science* **269**, 198–201 (1995); K. B. Davies, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, “Bose-Einstein condensation in a gas of sodium atoms”, *Phys. Rev. Lett.* **75**, 3969–73 (1995); C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, “Evidence of Bose-Einstein condensation in a atomic gas with attractive interactions”, *Phys. Rev. Lett.* **75**, 1687–90 (1995).
- [3] M. R. Andrews, C. G. Townsend, H. J. Miesner, D. S. Durfee, D. M. Kurn, and W. Ketterle, “Observation of interference between two Bose-Einstein condensates”, *Science* **275**, 637–41 (1997).
- [4] W. Ketterle and H. J. Miesner, “Coherence properties of Bose-Einstein condensates and atom lasers”, *Phys. Rev. A* **56**, 3291–3 (1997).
- [5] M. Lewenstein and L. You, “Quantum phase diffusion of a Bose-Einstein condensate”, *Phys. Rev. Lett.* **77**, 3489–93 (1996); M. J. Steel and M. J. Collett, “Quantum state of two trapped Bose-Einstein condensates with a Josephson coupling”, *Phys. Rev. A* **57**, 2920–30 (1998).
- [6] H. Zeng, W. Zhang, and F. Lin, “Non classical Bose-Einstein condensate”, *Phys. Rev. A* **52**, 2155–60 (1995).
- [7] J. Ruostekoski and J. Javanainen, “Quantum field theory of cooperative atom response: low light intensity”, *Phys. Rev. A* **55**, 513–26 (1997); E. A. Burt, R. W. Ghrist, C. J. Myatt, M. J. Holland, E. A. Cornell, and C. E. Wieman, “Coherence, correlations and collisions: what one learns about Bose-Einstein condensates from their decay”, *Phys. Rev. Lett.* **79**, 337–40 (1997).

- [8] See *e.g.* U. Leonhardt, *Measuring the Quantum State of Light* (Cambridge University Press, Cambridge, 1997); J. Mod. Opt. **44**, No. 11/12, Special issue *Quantum State Preparation and Measurement*.
- [9] S. Mancini and P. Tombesi, “Quantum state reconstruction of a Bose-Einstein condensate”, Europhys. Lett. **40**, 351–5 (1997).
- [10] E. L. Bolda, S. M. Tan, and D. F. Walls, “Reconstruction of the joint state of a two-mode Bose-Einstein condensate”, Phys. Rev. Lett. **79**, 4719–23 (1997); R. Walser, “Measuring the state of a bosonic two-mode quantum field”, Phys. Rev. Lett. **79**, 4724–27 (1997).
- [11] E. L. Bolda, S. M. Tan, and D. F. Walls, “Measuring the quantum state of a Bose-Einstein condensate”, Phys. Rev. A **57**, 4686–94 (1998).
- [12] D. T. Smithey, M. Beck, M. G. Raymer, and A. Faridani, “Measurement of the Wigner distribution and the density matrix of a light mode using optical homodyne tomography: application to squeezed states and the vacuum”, Phys. Rev. Lett. **70**, 1244–47 (1993).
- [13] R. Spreewitz, T. Pfau, U. Janicke, and M. Wilkens, “Laser-like scheme for atomic matter waves”, Europhys. Lett. **32**, 469–74 (1995); M. Holland, K. Burnett, C. Gardiner, J. I. Cirac, and P. Zoller, “Theory of an atom laser”, Phys. Rev. A **54**, 1757–60 (1996); H. Wiseman, A. M. Martins, and D. F. Walls, “An atom laser based on evaporative cooling”, Quant. Semiclass. Opt. **8**, 737–53 (1996); H. Steck, M. Naraschewski, and H. Wallis, “Output of a pulsed atom laser”, Phys. Rev. Lett. **80**, 1–5 (1998); J. Schneider and A. Schenzle, “Output from an atom laser: theory vs experiment” Appl. Phys. B **69**, 353–6 (1999); I. Bloch, T. W. Hänsch, and T. Esslinger, “Atom laser with a cw output coupler”, Phys. Rev. Lett. **82**, 3008–11 (1999).
- [14] J. A. Dunningham and K. Burnett, “Phase standard for Bose-Einstein condensates”, Phys. Rev. Lett. **82**, 3729–33 (1999).
- [15] S. Mancini, V. I. Man’ko, and P. Tombesi, “Different realizations of the tomography

- principle in quantum state measurement”, J. Mod. Opt. **44**, 2281–92 (1997).
- [16] G. M. D’Ariano, “Group theoretical quantum tomography”, Acta Physica Slovaca **49**, 513–22 (1999).
- [17] E. P. Wigner, *Perspectives in Quantum Theory*, W. Yourgrau and A. van der Merwe Eds., (Dover, New York, 1979).
- [18] *Atom Interferometry*, edited by P. Berman (Academic, New York, 1997); M.-O. Mewes, M. R. Andrews, D. M. Kurn, D. S. Durfee, C. G. Townsend, and W. Ketterle, “Output coupler for Bose-Einstein condensed atoms”, Phys. Rev. Lett. **78**, 582–5 (1997); H. Hinderthur, A. Pautz, F. Ruschewitz, K. Sengstock, and W. Ertmer, “Atomic interferometry with polarizing beam splitters”, Phys. Rev. A **57**, 4730–5 (1998).
- [19] *Quantum Theory of Angular Momentum*, L. C. Biedenharn and H. Van Dam, Eds. (Academic Press, New York, 1965).
- [20] V. V. Dodonov and V. I. Man’ko, “Positive distribution description for spin states”, Phys. Lett. A **229**, 335–9 (1997).
- [21] V. I. Man’ko and O. V. Man’ko, “Spin state tomography”, Zh. Eksp. Teor. Fiz. **112**, 796–800 (1997) [JETP **85**, 430–434 (1997)].
- [22] S. Wallentowitz and W. Vogel, “Unbalanced homodyning for quantum state measurements”, Phys. Rev. A **53**, 4528–33 (1996); K. Banaszek and K. Wodkiewicz, “Direct probing of quantum phase space by photon counting”, Phys. Rev. Lett. **76**, 4344–7 (1996); S. Mancini, P. Tombesi, and V. I. Man’ko, “Density matrix from photon number tomography”, Europhys. Lett. **37**, 79–83 (1997).
- [23] T. Opatrny and D. G. Welsch, “Density matrix reconstruction by unbalanced homodyning”, Phys. Rev. A **55**, 1462–5 (1997).
- [24] See *e.g.*: D. J. S. Robinson, *A course in Linear Algebra with Applications* (World Sci-

- entific, Singapore, 1991); W. T. Vetterling, S. A. Teukolsky, W. H. Press, and B. P. Flannery, *Numerical Recipes*, (Cambridge University Press, Cambridge, 1985).
- [25] M. O. Scully and W. E. Lamb, “Quantum theory of optical maser. III. Theory of photoelectron counting statistics”, Phys. Rev. **179**, 368–74 (1969).
- [26] K. Vogel and W. Schleich, in *Fundamental Systems in Quantum Optics*, J. Dalibard, J. M. Raimond and J. Zinn-Justin Eds., (Elsevier Science Publishers B. V., 1992).
- [27] M. Hillery, R. F. O’Connell, M. O. Scully, and E. P. Wigner, “Distribution functions in physics: Fundamentals”, Phys. Rep. **106**, 121–167 (1986). The Wigner function is a more refined picture of the quantum state in phase space, and is particularly suited to display quantum characters of the state. In simple words, the Q -function is a smoothed version of the Wigner function.
- [28] F. T. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, “Atomic coherent states in quantum optics” Phys. Rev. A **6**, 2211–37 (1972); G. S. Agarwal, “Relation between atomic coherent representation, state multipoles and generalized phase space distributions”, Phys. Rev. A **24**, 2889–96 (1981).
- [29] D. Forster, *Hydrodynamic Fluctuations, Broken Symmetry, and Correlation Functions*, (Benjamin, New York, 1975); K. Huang, *Statistical Mechanics* 2nd ed. (Wiley, New York, 1987).
- [30] G. M. D’Ariano, U. Leonhardt, and H. Paul, “Homodyne detection of the density matrix of the radiation field”, Phys. Rev. A **52**, R1801–4 (1995).
- [31] Proceedings of the Quark Matter ’96 Conference, (P. Braun-Munzinger, H. J. Specht, R. Stock, and H. Stoecker Eds.), Nucl. Phys. **A610**, 1c-565c (1996); Proceedings of the Strangeness in Hadronic Matter ’96 Conference, (T. Csorgo, P. Levai and J. Zimanyi Eds.), Heavy Ion Physics **4**, 1-440 (1996); S. Pratt, “Pion lasers from high energy collisions”, Phys. Lett. B **301**, 159-64 (1993).

FIGURES

FIG. 1. A squeezed 2-mode state. (a) Ideal Q -function when the displacement parameter is $x_0 = \sqrt{5}$ and the squeezing parameter is $r = e$. (b) The corresponding Q -function reconstructed through the method of Sec. III. To obtain this figure we have simulated experimental data by adding to each probability w a noise term with a Gaussian distribution, the latter having a width proportional to the ratio between the probability itself and the number of runs for given parameters.

FIG. 2. A squeezed state for the single mode. (a) Ideal Wigner function when the displacement parameter is $x_0 = \sqrt{3}$ and the squeezing parameter is $r = e$. (b) The corresponding Wigner function reconstructed through the method of Sec. IV. The reconstruction parameters are $|\beta| = 1.1$, $\eta = 0.9$, and 3×10^5 simulated experimental data per each phase have been used (see text).

FIG. 3. A number state for the single mode. (a) Ideal Wigner function for the Fock state $|n\rangle = |5\rangle$. (b) The corresponding Wigner function reconstructed through the method of Sec. IV. The reconstruction parameters are $|\beta| = 0.3$, $\eta = 0.9$, and 3×10^5 simulated experimental data per each phase have been used (see text).

FIG. 4. A squeezed state with a random phase between 0 and 2π . Here, a value of the displacement parameter $x_0 = \sqrt{5}$ has been used.

This figure "fig1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/quant-ph/0106089v1>

This figure "fig2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/quant-ph/0106089v1>

This figure "fig3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/quant-ph/0106089v1>

This figure "fig4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/quant-ph/0106089v1>